

Data Sheet

IBA Headquarters
IBA GmbH
Rudolf-Wissell-Str. 28
D-37079 Göttingen
Germany
Tel. +49 (0) 551-5 06 72-0
Fax +49 (0) 551-5 06 72-181
E-mail info@iba-go.com
<http://www.iba-go.com>

IBA US Distribution Center
10748 Indian Head Industrial
Blvd.
St. Louis, MO 63132
USA
Tel. 1-877-IBA-GmbH
(1-877-422-4624)
Fax 1-888-531-6813
E-mail info@iba-go.com
<http://www.iba-go.com>

pASK-IBA63c-plus

Cat. No. : 2-1463-300

Lot No.: 1463-

Last date of revision
May 10

Version 1463-300-3

Description	Expression plasmid. The expression cassette is under transcriptional control of the tetracycline promoter/operator.
Affinity tag	<i>Strep-Tactin</i> [®] affinity tag (<i>Strep-tag II</i> [®]) for the purification of recombinant protein. The affinity tag is fused to the C-terminus of the recombinant protein.
Bacterial Expression	Expression is induced upon addition of 200 µg anhydrotetracycline (order no.: 2-0401-001; 2-0401-002) per 1 liter <i>E. coli</i> shaking culture ($A_{550} = 0.5$).
Expression strain	Any <i>E. coli</i> strain. The <i>tet</i> -promoter works independently from the genetic background of <i>E. coli</i> .
Resistance	Ampicillin
Form	5 µg, dissolved in 10 mM Tris/HCl pH 8.0, 1 mM EDTA; 20 µl
Concentration	250 ng/µl
Storage	4 °C for frequent usage, -20 °C for long-term storage

For research use only

Important licensing information

This product is based on Strep-tag and tet promoter technologies covered by intellectual property (IP) rights and on completion of the sale IBA grants respective Limited Use Label Licenses to purchaser. IP rights and Limited Use Label Licenses for said technology are further described and identified at <http://www.iba-go.com/patents.html> or upon inquiry at info@iba-go.com or at IBA GmbH, Rudolf-Wissell-Str. 28, 37079 Göttingen, Germany. By use of this product the purchaser accepts the terms and conditions of all applicable Limited Use Label Licenses.

Trademark information


The owners of trademarks marked by "®" or "TM" are identified at <http://www.iba-go.com/patents.html>. Registered names, trademarks, etc. used in this document, even when not specifically marked as such, are not to be considered unprotected by law.

Multiple Cloning Site of pASK-IBA63c-plus

1	CCATCGAATGGCCAGATGATTAATTCCTAATTTTTGTTGACTCTATCATTGATAGAGTTATTTTACCACTCCCTA	77
	forward primer	
78	TCAGTGTAGAGAAAAGTGAAATGAATAGTTCGACAAAAATCTAGAAATAATTTTGTTTAACTTTAAGAAGGAGATATAC	157
	XbaI	
	<div style="display: flex; justify-content: space-around; font-size: small;"> M G Y L W I R I R A P S T S L R P H S R A L G A T R Strep-tag W S H P Q </div>	
158	CATGGGATATCTGTGGATCCGAATTCGAGCTCCGTCGACAAGCTTGCGGCCGCACTCGAGAGCGCTTGGAGCCACCCGCA	237
	NcoI EcoRV BamHI EcoRI SstI SalI HindIII NotI XhoI Eco47III	
	<div style="display: flex; justify-content: space-around; font-size: small;"> F E K * </div>	
238	GTTCGAAAAATAATGAGCTTGACCTGTGAAGTGAAAAATGGCGCACATTGTGCGACATTTTTTTGTCTGCCGTTTACCG	317
	BstBI reverse	
318	CTACTGCGTCACGGATCTCCACGCGCCCTGTAGCGCGCATTAAGCGCGCGGGTGTGGTTACGCGCAGCGTGACCG	397
	primer	

Features of pASK-IBA63c-plus

	from bp	to bp
promoter	37	72
forward primer binding site	57	76
multiple cloning site	157	223
Strep-tag	224	251
reverse primer binding site	309	325
f1 origin	338	776
AmpR resistance gene	925	1785
Tet-repressor	1795	2418
Col E1 origin	2571	3159



Sequencing primers:

Forward: 5' - GAGTTATTTTACCACTCCCT -3'

Reverse: 5' - CGCAGTAGCGGTAAACG -3'