

## Data Sheet

**IBA Headquarters**  
 IBA GmbH  
 Rudolf-Wissell-Str. 28  
 D-37079 Göttingen  
 Germany  
 Tel. +49 (0) 551-5 06 72-0  
 Fax +49 (0) 551-5 06 72-181  
 E-mail [info@iba-go.com](mailto:info@iba-go.com)  
<http://www.iba-go.com>

**IBA US Distribution Center**  
 10748 Indian Head Industrial Blvd.  
 St. Louis, MO 63132  
 USA  
 Tel. 1-877-IBA-GmbH  
 (1-877-422-4624)  
 Fax 1-888-531-6813  
 E-mail [info@iba-go.com](mailto:info@iba-go.com)  
<http://www.iba-go.com>

## pEXPR-IBA45

Cat. No. : 2-1945-000

Lot No.: 1945-

Last date of revision

March 06

Version 1945-8

<b>Description</b>	Eukaryotic expression vector designed for high-level stable and non-replicative transient expression in most mammalian hosts containing the following elements: <ul style="list-style-type: none"> <li>• Human cytomegalovirus (CMV) immediate-early promoter for high-level expression in a wide range of mammalian cells</li> <li>• Multiple cloning site</li> <li>• Neomycin resistance gene for selection of stable cell lines</li> <li>• Episomal replication in cell lines that are latently infected with SV40 or that express the SV40 large T antigen (e.g. COS-1, COS-7)</li> <li>• The expressed recombinant protein will be localized in the cytoplasm.</li> <li>• <i>Strep</i>-Tactin affinity tag (<i>Strep</i>-tag II) for the purification of recombinant protein via <i>Strep</i>-Tactin resins. The <i>Strep</i>-tag is fused to the N-terminus of the recombinant protein.</li> <li>• 6xHistidine-tag for the purification of recombinant protein via Ni-NTA resins. The 6xHistidine-tag is fused to the C-terminus of the recombinant protein.</li> </ul>
<b>Resistance for selection in <i>E. coli</i></b>	Ampicillin
<b>Form</b>	5 µg, dissolved in TE buffer (10 mM Tris/HCl pH 8.0, 1 mM EDTA); 20 µl
<b>Concentration</b>	250 ng/µl
<b>Storage</b>	4 °C for frequent usage, -20 °C for long-term storage
<b>Shipment</b>	Room temperature

### For research use only

*Strep*-tag® technology for protein purification and detection is covered by US patent 5,506,121, UK patent 2272698 and French patent 93 13 066; the tetracycline promoter based expression system is covered by US patent 5,849,576 and *Strep*-Tactin® is covered by US patent 6,103,493. Further patent applications are pending world-wide. Purchase of reagents related to these technologies from IBA provides a license for non-profit and in-house research use only. Expression or purification or other applications of above mentioned technologies for commercial use require a separate license from IBA. A license may be granted by IBA on a case-by-case basis, and is entirely at IBA's discretion. Please contact IBA for further information on licenses for commercial use. *Strep*-tag® and *Strep*-Tactin® are registered trademarks of IBA GmbH. Use of the CMV promoter is covered under U.S. patent nos. 5,168,062 and 5,385,839 owned and licensed by the University of Iowa Research Foundation and may be used for research purposes only. Commercial users must obtain a license to these patents directly from the University of Iowa Research Foundation. Inquiries for commercial use should be directed to: Brenda Akins, University of Iowa Research Foundation (UIRF), 214 Technology Innovation Center, Iowa City, IA 52242. The 6xHistidine-tag is licensed from Hoffmann-La Roche, Inc., Nutley, NJ and/or Hoffmann-La Roche Ltd., Basel, Switzerland and is provided only for the use in research. Information about licenses for commercial use is available from QIAGEN GmbH, Max-Volmer-Str. 4, D-40724 Hilden, Germany.


## Multiple Cloning Site of pEXPR-IBA45

721	AAAATCAACGGGACTTTCAAAAATGTCGTAACAACCTCCGCCCATTTACGCAAAATGGGCGGTAGGCGGTACGGTGGGAG	800
	CAAT	
801	GTCTATAAAGCAGAGCTCTCTGGCTAACTAGAGAACCACCTGCTTACTGGCTTATCGAAATTAATACGACTCACTATAG	880
	TATA <span style="margin-left: 150px;">forward primer</span>	
	<div style="display: flex; justify-content: space-around; width: 100%;"> <span>link</span> <span>Strep-tag</span> </div> <div style="display: flex; justify-content: space-around; width: 100%;"> <span>M</span> <span>A</span> <span>S</span> <span>W</span> <span>S</span> <span>H</span> <span>P</span> <span>Q</span> <span>F</span> <span>E</span> <span>K</span> <span>G</span> <span>A</span> <span>E</span> <span>T</span> <span>A</span> <span>V</span> <span>P</span> <span>N</span> <span>S</span> <span>S</span> <span>S</span> </div>	
881	GGTCTAGACCCACCATGGCTAGCTGGAGCCACCCGAGTTCGAAAAAGgcgCGAGACC GCGGTCCC GAATTCGAGCTCG	960
	<span>XbaI</span> <span style="margin-left: 100px;">NheI</span> <span style="margin-left: 100px;"><b>BbeI</b></span> <span style="margin-left: 20px;"><b>BsaI</b></span> <span style="margin-left: 100px;">EcoRI</span> <span style="margin-left: 50px;">KpnI</span>	
	<span style="margin-left: 100px;"><b>EheI</b></span> <span style="margin-left: 100px;">SacII</span>	
	<span style="margin-left: 100px;"><b>KasI</b></span> <span style="margin-left: 100px;"><b>NarI</b></span>	
	<div style="display: flex; justify-content: space-around; width: 100%;"> <span>link</span> <span>6xHistidine-tag</span> </div> <div style="display: flex; justify-content: space-around; width: 100%;"> <span>V</span> <span>P</span> <span>G</span> <span>D</span> <span>P</span> <span>S</span> <span>R</span> <span>S</span> <span>T</span> <span>C</span> <span>R</span> <span>G</span> <span>T</span> <span>M</span> <span>V</span> <span>S</span> <span>G</span> <span>L</span> <span>R</span> <span>G</span> <span>S</span> <span>H</span> <span>H</span> <span>H</span> <span>H</span> <span>H</span> <span>H</span> <span>*</span> </div>	
961	GTACCCGGGGATCCCTCGAGGTCGACCTGCAGGGGACCATGGTCTCAGgccTGAGAGGATCGCATCACCATCACCATCA	1040
	<span>BamHI</span> <span style="margin-left: 20px;">XhoI</span> <span style="margin-left: 100px;"><b>BsaI</b></span> <span style="margin-left: 20px;"><b>StuI</b></span>	
1041	CTAATAAGCTTGGCGCCGAGATCTAGCTTAAGTTTAAACCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCAT	1120
	<span>HindIII</span> <span style="margin-left: 100px;">PmeI</span> <span style="margin-left: 100px;">reverse primer</span>	
	NotI	

**Please note:** Restriction enzymes in bold cut twice, however, not all multiple cutting restriction enzymes are shown. The *BsaI* sites (isoschizomer of *Eco31I*) at each end of the multiple cloning site are useful for precise and oriented insertion of the recombinant gene by one cleavage reaction only. The “link” contains a restriction site which can be used e.g. for subcloning the recombinant gene into pASK-IBA vectors for bacterial expression.

## Features of pEXPR-IBA45

	from bp	to bp
CMV promoter	232	819
forward primer binding site	832	852
Strep-tag	904	927
multiple cloning site	928	1014
6xHistidine-tag	1015	1044
reverse primer binding site	1091	1108
f1 origin	1367	1795
Neomycin resistance gene	2205	2999
pUC origin	3686	4356
Ampicillin resistance gene	4501	5361



Cloning primers for the precise cloning using <i>BsaI</i> or <i>Eco31I</i>	Sequencing primers:
Forward: 5'- NNNNNNGGTCTCNGC GCC <sup>(N<sub>20</sub>)</sup> NNN NNN ...	Forward: 5'- GAGAACCACCTGCTTACTGGC -3'
Reverse: 5'- NNNNNNGGTCTCNG GCC <sup>(N<sub>20</sub>)</sup> NNN NNN ...	Reverse: 5'- TAGAAGGCACAGTCGAGG -3'